

Performance of Modified Pneumonia Severity Indexes in Community-Acquired Pneumonia

Yasemin Yılmaz¹, Aynur Yurtseven²

¹University of Health Sciences Türkiye, Ankara Atatürk Sanatorium Training and Research Hospital, Clinic of Emergency Medicine, Ankara, Türkiye

²University of Health Sciences Türkiye, Etlik City Hospital, Clinic of Emergency Medicine, Ankara, Türkiye

Abstract

Aim: In patients with community-acquired pneumonia, we observed that the impact of the age variable commonly used in Pneumonia Severity scores to predict mortality risk is disproportionate within the pneumonia severity index (PSI). In this context, our study aimed to evaluate and compare the effectiveness of pneumonia severity scores, adjusted for demographic factors such as age and gender, in predicting intensive care unit (ICU) admission and mortality, compared to PSI and CURB-65.

Materials and Methods: In 2019, data on patients diagnosed with pneumonia in the emergency department were obtained through a retrospective review. The CURB-65, PSI, age-modified PSI (mPSI), and gender-modified PSI (gmPSI) scores were calculated. The predictive performance of these scores for ICU admission and mortality was statistically analyzed.

Results: A total of 363 patients were included in the study, of whom 205 (56.4%) were male. Additionally, comparisons of the newly developed age-mPSI and gmPSI with the standard PSI and CURB-65 demonstrated no statistically significant difference in predicting 30-day mortality or ICU admission rates ($p>0.05$).

Conclusion: The findings of this study indicate that the newly developed mPSI and gmPSI scoring systems yield results comparable to those of the standard PSI scoring system.

Keywords: Community acquired pneumonia, pneumonia severity index, CURB-65, age, gender

Introduction

Community-acquired pneumonia (CAP) is a common disease with a lifetime prevalence of 20-30% in developing countries and 3-4% in developed countries (1). The mortality rate among hospitalized and diagnosed patients has been reported to vary between 4% and 21% in different settings (2). It has been reported that this rate is higher, even up to 50%, among patients admitted to the intensive care unit (ICU) (3). Treatment in patients with CAP is determined according to the general condition and prognosis of the patient, and the follow-up and treatment plan of the patient varies according to the severity of the disease (4). Therefore, various scoring systems are used to determine the prognosis. The most commonly used assessment tools are CURB-65 (confusion, blood urea nitrogen, respiratory rate, blood pressure) and

Pneumonia Severity index (PSI). CURB-65 is a classification system used to predict the risk of mortality and is frequently preferred in emergency departments because it is easy to remember and contains few criteria (3). PSI is a scoring system developed by Fine et al. (5) that includes 20 parameters and is divided into 5 categories according to 30-day prognosis predictions. CURB-65 and the PSI are widely used to predict mortality in CAP (6). However, there are many studies reporting that PSI performs better than CURB-65 in predicting mortality (7,8).

Nevertheless, clinical applications have revealed certain limitations of this scoring system within the context of emergency departments (8,9). For instance, one study demonstrated that mortality and ICU admission rates were lower among patients categorized as PSI Class I/II but were paradoxically higher among those with a CURB-65 score of 1 (10). Furthermore, evidence



Corresponding Author: Yasemin Yılmaz MD, University of Health Sciences Türkiye, Ankara Atatürk Sanatorium Training and Research Hospital, Clinic of Emergency Medicine, Ankara, Türkiye
E-mail: ysmnyaydin@gmail.com **ORCID ID:** orcid.org/0000-0003-1048-4478

Cite this article as: Yılmaz Y, Yurtseven A. Performance of modified pneumonia severity indexes in community-acquired pneumonia. Eurasian J Emerg Med. [Epub Ahead of Print].

Received: 11.02.2025

Accepted: 09.05.2025

Epub: 07.07.2025



©Copyright 2025 The Emergency Physicians Association of Turkey / Eurasian Journal of Emergency Medicine published by Galenos Publishing House.
Licenced by Creative Commons Attribution-NonCommercial-NoDerivatives (CC BY-NC-ND) 4.0 International License.

suggests that approximately 27% of patients in PSI Class I-III may necessitate ICU admission and 40% of patients classified as low-risk were hospitalized (11,12). Although elevated scores in such systems are typically indicative of more severe illness, it is important to note that younger to middle-aged patients with low scores may also require ICU-level care. In particular, the age-related risk factor incorporated into the PSI scoring system may result in disproportionately high scores among older patients while potentially underestimating severity in younger individuals.

Many studies in the literature have shown that sex hormones play an important role in the immune response and that women have a stronger immune response than men. In general, women have a stronger response to pathogens and produce higher levels of interferon and antibodies (13,14). We aimed to assess the influence of age and sex on the performance of the PSI scoring system in this study. In particular, the impact of the newly developed age- and sex-modified, modified-PSI (mPSI) and gender- modified PSI (gmPSI) on clinical outcomes, including mortality and ICU admission rates, was analyzed.

Materials and Methods

This retrospective study was conducted in the emergency department of a tertiary hospital located in a provincial center that experiences approximately 420,000 patient visits annually. It received ethical approval from the University of Health Sciences Türkiye, Dışkapı Yıldırım Beyazıt Training and Research Hospital Clinical Research Ethics Committee (decision number: 103/13, date: 25.01.2021). Our research was designed in accordance with the Standards for Reporting of Diagnostic Accuracy Studies statement (15).

Information was extracted from electronic medical records and patient charts. A retrospective chart review was conducted by two emergency medicine specialists, each possessing a minimum of three years of experience.

Between January 1, 2019, and December 31, 2019, patients were diagnosed with pneumonia in the emergency medicine clinic. All patients over the age of 18 years with a diagnosis of pneumonia were included in the study. The inclusion criteria were based on ICD-10 codes (International Statistical Classification of Diseases, J15 and its subcodes, J15.8, J15.9) retrieved from the hospital's data system. Individuals with a history of hospitalization within the preceding seven days, a diagnosis of cystic fibrosis, tuberculosis, or Human immunodeficiency virus, a documented history of concomitant pulmonary embolism, or a history of trauma at the time of admission, as well as those with incomplete data in their records, were excluded from the study. Additionally,

patients with high mortality at the time of admission and those diagnosed with pneumonia were not included in the study.

Patients diagnosed with pneumonia were initially stratified according to the CURB-65 score, the PSI, and the criteria for ICU admission (1). The PSI scoring system was subsequently modified by excluding the age and gender variables. To adjust for age-related risk, 10 points were allocated to patients aged 65 years and older, whereas no points were assigned to those under 65 years. This revised scoring model was termed the mPSI.

In the calculation of the gmPSI, female patients aged 65 years and older were assigned 10 points, while male patients in the same age group were allocated 20 points. No points were assigned to either male or female patients under the age of 65.

The predictive performance of the scoring systems (PSI, mPSI, and gmPSI) in estimating hospitalization and mortality outcomes was systematically evaluated. Patients were subsequently categorized into two distinct age groups: younger adults (18-64 years) and older adults (≥ 65 years), enabling a comprehensive comparison between these cohorts.

Statistical Analysis

Statistical analyses were conducted using the SPSS 22.0 software package. The normality of continuous variables was assessed through the Kolmogorov-Smirnov test, supplemented by descriptive statistical measures of Skewness and Kurtosis.

Descriptive statistics were presented as frequencies and percentages for categorical variables, while continuous variables were summarized using medians and (interquartile range 25th-75th percentiles). For the analysis of differences in numerical variables between two independent groups, the Mann-Whitney U test was employed. Comparisons of proportions between independent groups were conducted using chi-square analysis.

ROC analyses were conducted to evaluate the predictive efficacy of continuous variables in estimating 30-day mortality and the need for intensive care. A p-value of <0.05 was considered the threshold for statistical significance.

Results

A total of 450 patients were initially enrolled in the study; however, 87 were excluded due to incomplete data. Among the remaining participants, 205 (56.4%) were male. Hospitalization rates did not differ significantly between male and female patients ($p=0.248$).

The stratified pneumonia scores based on 30-day mortality and ICU requirements for all patients are presented in Table 1. Patients who died or required ICU admission were older and

exhibited higher scores across all measurements ($p < 0.05$ for all comparisons) (Table 1).

ROC analysis was conducted to identify the threshold values of pneumonia scores distinguishing between deceased and surviving patient groups. Among the scores, PSI demonstrated the highest area under the curve (AUC), while CURB-65 exhibited the lowest. All AUC values were statistically significant ($p < 0.001$ for all comparisons) (Table 2, Figure 1A).

In the ROC analysis conducted to determine the threshold values of pneumonia scores differentiating patients admitted to the ICU from those not admitted, the mPSI demonstrated the highest AUC, while the CURB-65 showed the lowest. All AUC values were statistically significant ($p < 0.001$ for all comparisons) (Table 3, Figure 1B).

The best cut-off values, along with the corresponding sensitivity and specificity of the PSI, mPSI, and gmPSI scores for predicting mortality and ICU admission, are presented in Table 4. The sensitivity and specificity values for all three scoring systems were found to be comparable (Table 4).

Patients were categorized into two age groups: 18-64 years (young) and ≥ 65 years (geriatric). In the ROC analysis conducted to identify the threshold values of pneumonia scores for predicting mortality within the young age group, the PSI demonstrated the highest AUC, whereas CURB-65 exhibited the lowest. All AUC values were statistically significant ($p < 0.001$) (Table 5, Figure 2A).

In the ROC analysis conducted to determine the threshold values of pneumonia scores for predicting mortality in the geriatric group, PSI and mPSI exhibited the highest AUC values, while CURB-65 demonstrated the lowest (Table 6, Figure 2B).

In the ROC analysis conducted to identify the threshold values of pneumonia scores for predicting ICU admission in the young patient group, the highest AUC value was observed for the mPSI, while the lowest was associated with the PSI (Table 7, Figure 3A).

In the ROC analysis conducted to determine the threshold values of pneumonia scores for intensive care admission in the geriatric population, the highest AUC value was observed for the PSI score, while the lowest was associated with the CURB-65 score (Table 8, Figure 3B).

Discussion

Treatment, prognosis, and care location decisions are often facilitated by scoring systems, which, despite inherent challenges and limitations, serve as valuable tools in the clinical decision-making process. These systems primarily aim to minimize unnecessary hospital admissions while ensuring appropriate patient placement and management. The PSI, a widely utilized scoring system, comprises 20 variables encompassing patient age, history of nursing home residence, comorbid conditions, vital signs, laboratory results, chest radiographic findings, and oxygenation status. The assessment is conducted in two distinct stages. In this scoring system, the age variable is directly assigned based on the numerical age in male patients, while, for female patients, it is calculated as age minus 10. Consequently, this approach may result in low scores for young patients with severe respiratory failure in the absence of comorbidities, while generating high scores for very elderly patients. Such discrepancies can create confusion for clinicians and potentially lead to inappropriate hospitalization and treatment decisions.

Table 2. AUC values of pneumonia scores according to 30-day mortality

	AUC	Confidence interval	p-value
CURB-65	0.700	0.637-0.763	<0.001
PSI	0.803	0.750-0.855	<0.001
mPSI	0.794	0.740-0.849	<0.001
gmPSI	0.793	0.739-0.847	<0.001

AUC: Area under the curve, PSI: Pneumonia severity index, mPSI: Modified-PSI, gmPSI: Gender-mPSI

Table 1. Pneumonia severity scores stratified by 30-day mortality and intensive care unit admission requirement

	Mortality in 30 days			ICU Hospitalization		
	Exitus	Alive	p-value	None ICU admission	ICU admission	p-value
Gender n (%)						
Female	33 (44.6%)	125 (43.3%)	0.835	106 (41.4%)	52 (48.6%)	0.248
Age (years) (min-max)	80 (70-86)	75 (63-83.5)	0.005	75 (63-81.7)	80 (70-87)	<0.001
CURB-65 median (min-max)	2 (2-3)	2 (1-2)	<0.001	2 (1-2)	2 (2-3)	<0.001
PSI score	164 (138-191)	119 (93-149)	<0.001	112.5 (91-140)	163 (142-180)	<0.001
mPSI, median (min-max)	100 (80-120)	60 (40-80)	<0.001	50 (30-80)	100 (80-120)	<0.001
gmPSI, median (min-max)	100 (80-120)	60 (40-90)	<0.001	60 (40-80)	100 (90-120)	<0.001
ICU admission n (%)	55 (74.3%)	58 (20.1%)	<0.001	-	-	-

ICU: Intensive care unit, PSI: Pneumonia severity index, mPSI: Modified-PSI, gmPSI: Gender-mPSI

The literature contains numerous studies evaluating the sensitivity and specificity of CURB-65 and PSI scores in predicting mortality and the necessity for ICU admission (16-18). In a study conducted in Iran, CURB-65 demonstrated higher accuracy in predicting mortality and the need for intensive care hospitalization in patients with CAP (17). In contrast, studies conducted by Akça et al. (17) and Satıcı et al. (18) found that PSI exhibited superior performance in predicting mortality.




Figure 1. ROC analysis of pneumonia scores according to 30-day mortality or ICU hospitalization for all patients. A) ROC analysis of pneumonia scores according to 30-day mortality for all patients, B) ROC analysis of pneumonia scores according to ICU hospitalization for all patients

ICU: Intensive care unit

	AUC	Confidence interval	p-value
CURB-65	0.764	0.714-0.815	<0.001
PSI	0.839	0.796-0.882	<0.001
mPSI	0.856	0.818-0.895	<0.001
gmPSI	0.854	0.815-0.892	<0.001

AUC: Area under the curve, ICU: Intensive care unit, PSI: Pneumonia severity index, mPSI: Modified-PSI, gmPSI: Gender-mPSI

	PSI	mPSI	gmPSI
Mortality			
Best cut-off point	121%	57%	52%
Sensitivity	89%	90%	95%
Specificity	55%	47%	42%
ICU			
Best cut-off point	123%	62%	62%
Sensitivity	90%	92%	94%
Specificity	62%	66%	62%

ICU: Intensive care unit, PSI: Pneumonia severity index, mPSI: Modified-PSI, gmPSI: Gender-mPSI

In a retrospective study involving 1,902 patients, Zhang et al. (19) categorized the patients into three age groups: 18-64 years, 65-84 years, and ≥ 85 years. Although PSI Class III is considered a low-risk category for mortality, they observed a mortality rate of 7.5% in the 18-64 age group, compared to 2.1% in the 65-84 age group and 2.6% in those aged ≥ 85 years. Accordingly, they suggested that PSI Class III may not be appropriate for classification as a low mortality risk in younger patients. They reported that while

Table 5. AUC values of pneumonia scores according to 30-day mortality at 18-64 years of age

	AUC	Confidence interval	p-value
CURB-65	0.760	0.594-0.927	0.008
PSI	0.870	0.756-0.985	<0.001
mPSI	0.825	0.688-0.963	=0.001
gmPSI	0.825	0.687-0.962	=0.001

AUC: Area under the curve, PSI: Pneumonia severity index, mPSI: Modified-PSI, gmPSI: Gender-mPSI




Figure 2. ROC analysis of pneumonia scores according to 30-day mortality for young and geriatric patients. A) ROC analysis of pneumonia scores according to 30-day mortality for young patients, B) ROC analysis of pneumonia scores according to 30-day mortality for geriatric patients




Figure 3. ROC analysis of pneumonia scores according to ICU hospitalization for young and geriatric patients. A) ROC analysis of pneumonia scores according to ICU hospitalization for young patients, B) ROC analysis of pneumonia scores according to ICU hospitalization for geriatric patients

ICU: Intensive care unit

Table 6. AUC values of pneumonia scores for predicting 30-day mortality in patients aged ≥65 years

	AUC	Confidence interval	p-value
CURB-65	0.658	0.585-0.731	<0.001
PSI	0.771	0.709-0.834	<0.001
mPSI	0.771	0.708-0.834	<0.001
gmPSI	0.769	0.707-0.832	<0.001
AUC: Area under the curve, PSI: Pneumonia severity index, mPSI: Modified-PSI, gmPSI: Gender-mPSI			

Table 7. AUC values of pneumonia scores for 18-64 years according to ICU

	AUC	Confidence interval	p-value
CURB-65	0.865	0.775-0.955	<0.001
PSI	0.835	0.726-0.944	<0.001
mPSI	0.901	0.814-0.988	<0.001
gmPSI	0.900	0.814-0.987	<0.001
AUC: Area under the curve, ICU: Intensive care unit, PSI: Pneumonia severity index, mPSI: Modified-PSI, gmPSI: Gender-mPSI			

Table 8. AUC values of pneumonia scores for predicting intensive care unit admission in the geriatric population

	AUC	Confidence interval	p-value
CURB-65	0.728	0.666-0.790	<0.001
PSI	0.833	0.787-0.880	<0.001
mPSI	0.830	0.783-0.876	<0.001
gmPSI	0.828	0.781-0.875	<0.001
AUC: Area under the curve, ICU: Intensive care unit, PSI: Pneumonia severity index, mPSI: Modified-PSI, gmPSI: Gender-mPSI			

PSI was more sensitive than CURB-65 in predicting mortality, it misclassified the low-risk group among younger patients, and the discriminative power of both CURB-65 and PSI diminished with increasing age (19). In a prospective study involving 987 patients, Chen et al. (20) categorized the patients into age groups as defined in the study and observed mortality rates of 20% in the 18-64 age group, 11% in the 65-84 age group, and 11.5% among those aged 85 years and over. They demonstrated that 77.8% of the patients who died in the young adult group were categorized as low-risk, a significantly higher proportion compared to those categorized as low-risk in the elderly and very old groups. They attributed the poor performance of the PSI score in elderly patients to the significant influence of the age variable and proposed that applying a non-linear adjustment to the age variable would enhance the PSI score's accuracy. Simonetti et al. (21) also observed in their prospective study that the effectiveness of PSI and CURB-65 in predicting mortality diminished with advancing age. In our clinical practice, we found

it necessary to critically evaluate the age component of the PSI score. Our observations indicated that some very elderly patients did not require intensive care despite having a high PSI score, whereas certain younger patients required intensive care despite a low PSI score.

In our study, a novel scoring system was developed and implemented to address the confusion caused by existing scoring systems for patients requiring hospitalization in the ward or ICU, or those who can be managed as outpatients. Based on the data obtained from these scoring systems, we found that the highest AUC value for predicting 30-day mortality was associated with PSI, while the lowest was observed with CURB-65.

Comparisons according to the patients' age groups indicated that the highest AUC value for 30-day mortality in the 18-64 age group was related to PSI, and the highest AUC value for the need for ICU admission was related to mPSI. We found that the highest AUC value for 30-day mortality in the geriatric age group was related to both PSI and mPSI, and the highest AUC value for the need for ICU admission was related to PSI. PSI scoring was more significant than mPSI scoring in terms of the need for intensive care admission in the geriatric age group. However, the results of both scores were similar in the geriatric age group in terms of mortality. In the 18-64 age group, the significance of mPSI scoring in intensive care admissions led to the conclusion that the age factor had an effect on determining the need for admission. However, after a more detailed examination of the findings, we found that there was no significant difference between the scoring systems.

Study Limitations

Our study has a retrospective design, and the consequent data loss is one of our limitations. There may be missing data due to incorrect diagnosis code records. Due to the retrospective nature of the study, certain variables like confusion may have been under-documented in the medical records, which could have resulted in lower CURB-65 and PSI scores. There are limitations in our study because patient data were collected in a single center and the study was conducted with a specific sample size.

Conclusion

In our study, we have demonstrated that the mPSI and gmPSI scoring systems, can reduce hospital burden and additional costs by decreasing unnecessary intensive care admissions, similar to PSI. Additionally, they exhibit comparable sensitivity to PSI, which is used as the standard for predicting early mortality. Therefore, we believe that its application in both elderly and younger patients could facilitate a more objective assessment.

Ethics

Ethics Committee Approval: It received ethical approval from the University of Health Sciences Türkiye, Dışkapı Yıldırım Beyazıt Training and Research Hospital Clinical Research Ethics Committee (decision number: 103/13, date: 25.01.2021).

Informed Consent: Retrospective study.

Footnotes

Authorship Contributions

Surgical and Medical Practices: Y.Y., A.Y., Concept: Y.Y., A.Y., Design: Y.Y., A.Y., Data Collection or Processing: Y.Y., A.Y., Analysis or Interpretation: Y.Y., A.Y., Literature Search: Y.Y., A.Y., Writing: Y.Y.

Conflict of Interest: The authors declare that they have no conflicts of interest.

Financial Disclosure: There are no financial conflicts of interest to disclose.

References

- Mandell LA, Wunderink RG, Anzueto A, Bartlett JG, Campbell GD, Dean NC, et al. Infectious Diseases Society of America/American thoracic society consensus guidelines on the management of community-acquired pneumonia in adults. *Clin Infect Dis*. 2007;44 Suppl 2:S27-72.
- Shah BA, Ahmed W, Dhobi GN, Shah NN, Khursheed SQ, Haq I. Validity of pneumonia severity index and CURB-65 severity scoring systems in community acquired pneumonia in an Indian setting. *Indian J Chest Dis Allied Sci*. 2010;52:9-17.
- Lim WS, van der Eerden MM, Laing R, Boersma WG, Karalus N, Town GI, et al. Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation study. *Thorax*. 2003;58:377-82.
- Kara PH, Demircan A, Akinçi E, Bildik F, Aygencel G, Ozsarac M. Focal mass-like opacity on chest radiography: round pneumonia. *J Emerg Med*. 2010;39:e89-90.
- Fine MJ, Auble TE, Yealy DM, Hanusa BH, Weissfeld LA, Singer DE, et al. A prediction rule to identify low-risk patients with community-acquired pneumonia. *N Engl J Med*. 1997;336:243-50.
- Oktariani Nur R, Pitoyo Ceva W, Singh Gurmeetb, Mansjoer Arifa. CURB 65 score as a predictor of early mortality in hospital-acquired pneumonia. *The Egyptian Journal of Chest Diseases and Tuberculosis*. 2019;68:231-5.
- Aujesky D, Auble TE, Yealy DM, Stone RA, Obrosky DS, Meehan TP, et al. Prospective comparison of three validated prediction rules for prognosis in community-acquired pneumonia. *Am J Med*. 2005;118:384-92.
- İlhan G, Karakaya Z, Akyol PY, Topal FE, Payza U. Evaluation of PSI and CURB-65 scoring systems which patients were diagnosed as community-acquired pneumonia in emergency department. *CMJ*. 2017;39:586-96.
- Olçay HÖ, Çorbacıoğlu ŞK, Emektar E, Çevik Y. Effectiveness of the YEARS algorithm added to classical clinical decision-making rules in suspected pulmonary embolism. *Glob Emerg Crit Care*. 2024;3:58-62.
- Ananda-Rajah MR, Charles PG, Melvani S, Burrell LL, Johnson PD, Grayson ML. Comparing the pneumonia severity index with CURB-65 in patients admitted with community acquired pneumonia. *Scand J Infect Dis*. 2008;40:293-300.
- Rosón B, Carratalà J, Dorca J, Casanova A, Manresa F, Gudiol F. Etiology, reasons for hospitalization, risk classes, and outcomes of community-acquired pneumonia in patients hospitalized on the basis of conventional admission criteria. *Clin Infect Dis*. 2001;33:158-65.
- Angus DC, Marrie TJ, Obrosky DS, Clermont G, Dremsizov TT, Coley C, et al. Severe community-acquired pneumonia: use of intensive care services and evaluation of American and British Thoracic Society Diagnostic criteria. *Am J Respir Crit Care Med*. 2002;166:717-23.
- Taneja V. Sex hormones determine immune response. *Front Immunol*. 2018;9:1931.
- Straub RH. The complex role of estrogens in inflammation. *Endocr Rev*. 2007;28:521-74.
- Cohen JF, Korevaar DA, Altman DG, Bruns DE, Gatsonis CA, Hooft L, et al. STARD 2015 guidelines for reporting diagnostic accuracy studies: explanation and elaboration. *BMJ Open*. 2016;6:e012799.
- Alavi-Moghaddam M, Bakhshi H, Rezaei B, Khashayar P. Pneumonia severity index compared to CURB-65 in predicting the outcome of community acquired pneumonia among patients referred to an Iranian emergency department: a prospective survey. *Braz J Infect Dis*. 2013;17:179-83.
- Akça HŞ, Algin A, Özdemir S, Sevimli H, Kokulu K, Eroğlu SE. Comparison of the efficacy of PSI, CURB-65, CALL and BCRSS in predicting prognosis and mortality in COVID-19 patients. *J Exp Clin Med*. 2021;38:434-9.
- Satici C, Demirkol MA, Sargin Altunok E, Gursoy B, Alkan M, Kamat S, et al. Performance of pneumonia severity index and CURB-65 in predicting 30-day mortality in patients with COVID-19. *Int J Infect Dis*. 2020;98:84-9.
- Zhang ZX, Yong Y, Tan WC, Shen L, Ng HS, Fong KY. Prognostic factors for mortality due to pneumonia among adults from different age groups in Singapore and mortality predictions based on PSI and CURB-65. *Singapore Med J*. 2018;59:190-8.
- Chen JH, Chang SS, Liu JJ, Chan RC, Wu JY, Wang WC, et al. Comparison of clinical characteristics and performance of pneumonia severity score and CURB-65 among younger adults, elderly and very old subjects. *Thorax*. 2010;65:971-7.
- Simonetti AF, Viasus D, Garcia-Vidal C, Carratalà J. Management of community-acquired pneumonia in older adults. *Ther Adv Infect Dis*. 2014;2:3-16.